



ORIGINAL RESEARCH

Sociodemographic factors associated with knowledge of type 2 diabetes in rural Tamil Nadu, India

AUTHORS



Hana MacDonald¹ MPH, Data Analysis Coordinator, Durham Region



Andrew Papadopoulos² PhD, Associate Professor



Cate Dewey³ PhD, Associate Professor



Sally Humphries⁴ PhD, Professor Emerita



Warren Dodd⁵ PhD, Assistant Professor



Kirit Patel⁶ PhD, Associate Professor



Matthew Little⁷ PhD, Assistant Professor *

CORRESPONDENCE

*Dr Matthew Little matthewlittle@uvic.ca

AFFILIATIONS

^{1,2} Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada

³ Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada; and Centre for Public Health and Zoonoses, Guelph, ON, Canada

⁴ Department of Sociology and Anthropology, University of Guelph, Guelph, ON, Canada

⁵ Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada; and School of Public Health Sciences, University of Waterloo, Waterloo, ON, Canada

⁶ International Development Studies, Menno Simons College, Canadian Mennonite University, Winnipeg, MB, Canada

⁷ Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada; and School of Public Health and Social Policy, University of Victoria, Victoria, BC, Canada

PUBLISHED

20 January 2022 Volume 22 Issue 1

HISTORY

RECEIVED: 27 April 2021

REVISED: 12 October 2021

ACCEPTED: 19 October 2021

CITATION

MacDonald H, Papadopoulos A, Dewey C, Humphries S, Dodd W, Patel K, Little M. Sociodemographic factors associated with knowledge of type 2 diabetes in rural Tamil Nadu, India. *Rural and Remote Health* 2022; 22: 6855. <https://doi.org/10.22605/RRH6855>

This work is licensed under a [Creative Commons Attribution 4.0 International Licence](https://creativecommons.org/licenses/by/4.0/)

ABSTRACT:

Introduction: This study aimed to investigate awareness of type 2 diabetes and how sociodemographic factors influence diabetes knowledge in a rural population of Tamil Nadu, India. Previous research has identified poor awareness of diabetes in several low and middle-income countries, which can lead to a high prevalence of undiagnosed diabetes. India having the second highest prevalence of diabetes globally, it is increasingly important to assess how diabetes can be addressed in rural Indian populations.

Methods: Systematic random sampling was used to gather study participants in 17 villages within the Krishnagiri district of Tamil Nadu, India. Data on diabetes knowledge was collected using a validated questionnaire. Knowledge score range was 0–8; a score of zero was designated as 'low knowledge', scores 1–4 as 'moderate knowledge', and scores 5–8 as 'good knowledge'. Associations between sociodemographic factors and composite diabetes knowledge score were assessed using a multinomial logistic GLLAMM model in Stata.

Results: A total of 753 individuals participated in the study. The average age of participants was 47 years and 55% were women. Overall awareness of diabetes was low, with 66% of individuals having no knowledge of diabetes. Only 16% and 17% achieved a moderate and a good knowledge score, respectively. Achieving a

moderate knowledge score was significantly positively associated with education, wealth, participation in the *Mahatma Gandhi National Rural Employment Guarantee Act* (MGNREGA), and business ownership as a source of income. Achieving a good knowledge score was significantly positively associated with education, wealth, rurality, participation in MGNREGA, business ownership as a source of income, and frequency of healthcare utilization. Rurality was significantly negatively associated (relative risk ratio (95% confidence interval)) with both moderate knowledge score (0.34 (0.19–0.59)), and good knowledge score (0.43 (0.24–0.74)). The strongest predictor of having a good knowledge score was having a high-school graduate or post-secondary education (11.07 (4.44–27.61)). Enrolment in MGNREGA employment was the strongest predictor for having a moderate knowledge score (3.27 (1.93–5.54)), as well as strongly associated with having a good knowledge score (2.39 (1.31–4.36)).

Conclusion: The low awareness of diabetes among participants of this study raises serious concerns for public health in India. Public health efforts must prioritize health equity to lessen the impacts of diabetes in rural populations, where individuals face systemic barriers to receiving prevention and treatment for conditions such as diabetes.

Keywords:

diabetes awareness, health education, health promotion, India, type 2 diabetes.

FULL ARTICLE:

Introduction

Type 2 diabetes mellitus (hereafter 'diabetes') is currently one of the fastest growing health emergencies in the world. The International Diabetes Federation reported that diabetes caused approximately four million deaths in 2019¹. India has experienced a large increase in non-communicable diseases in the past several decades², and now has the second greatest number of people with diabetes in the world at 77 million individuals¹. This number is expected to increase to over 101 million people with diabetes in the next 10 years¹. Broader societal processes such as urbanization and globalization, along with dietary changes and declining physical activity, have been primary drivers of the diabetes epidemic in low- and middle-income countries such as China and

India³. Risk of developing diabetes is dependent on both modifiable factors (eg lack of physical activity and obesity) and non-modifiable factors (eg age and family history of diabetes)⁴. Diabetes carries a large economic burden due to disability and death from several complications, including retinopathy, kidney failure, and heart disease, among others⁵. The long-lasting pathology of disease complications can have severe impacts on individuals, families, communities, and healthcare systems. Individuals with diabetes from families of low socioeconomic status, who already face financial hardships, often spend as much as 60% of their household income to care for family members with diabetes³.

Many studies have shown that diabetes is an increasing concern in

South India, and especially in the state of Tamil Nadu^{2,6,7}. Diabetes affects approximately 4.8 million individuals in Tamil Nadu⁸, with a current age-standardized prevalence of 10.8%⁷. The increasing prevalence of diabetes in Tamil Nadu has been associated with increased waist-to-hip ratio, BMI, tobacco consumption, urban residency, decreased physical activity⁹, as well as male sex, age, obesity, scheduled castes, family history of diabetes^{4,10}, alcohol intake, and hypertension¹¹.

The low awareness of diabetes and prediabetes in India has also been highlighted in several studies¹²⁻¹⁷. Low awareness can lead to high prevalence of undiagnosed diabetes, which is a serious public health concern; indeed, estimates suggest that approximately half of diabetes cases are undiagnosed in India¹⁸. It is therefore necessary to investigate both the level of awareness of, as well as factors associated with knowledge of, diabetes, to identify priority areas for targeting educational campaigns, public health surveillance, and diabetes screening. This is particularly true in rural and socioeconomically disadvantaged regions, where limited access to education and healthcare resources is a substantial barrier to health literacy and diabetes awareness^{19,20}. Within this context, the primary objectives of this study were to assess the overall awareness of diabetes, and to determine associations between sociodemographic factors and diabetes knowledge in a rural population located in north-western Tamil Nadu. With a high statewide prevalence of diabetes⁷, it is increasingly important to ensure relevant health education is targeted towards high-risk groups in Tamil Nadu to mitigate the ongoing diabetes epidemic.

Methods

Study design and data collection

Data were collected as a component of a cross-sectional health study conducted in 17 villages in a rural region of Krishnagiri District, Tamil Nadu⁹. Previous publications using these data have identified and discussed other specific health outcomes from this study, including newly diagnosed diabetes⁹, overweight and obesity²¹, and anemia²²; however, this analysis is the first conducted on diabetes awareness using this dataset. Specific sampling techniques, including a sample size calculation, as well as other data collection methods, are described in detail elsewhere⁹. In brief, the research team conducted systematic random sampling to recruit adult participants (≥ 19 years) using the Kish method²³. Following recruitment and informed consent, a survey was administered to participants by a trained researcher to collect information on demographics, occupation and livelihood characteristics, self-reported health, and household assets. Knowledge on diabetes was assessed using a validated questionnaire developed by Mohan et al (2005), which consisted of five questions assessing awareness of diabetes, including symptoms, causes, and complications²⁴.

Definitions and explanations of variables

Although all villages included in the study were rural by the definition as described by the Census of India, the rurality of each

village was assessed as a predictor variable using a rurality index, adapted from Weinert and Boik²⁵. The two variables incorporated into the rurality index were distance to the primary healthcare center (given half a positive weighted value), and the population size of each village (given a full negative weighted value). The results were standardized to a mean of zero and standard deviation of one, with a positive score indicating a more rural residence, and a negative score reflecting a less rural residence.

Data on household assets and community facilities (such as toilet facilities and water pumps, if applicable) were collected to assess socioeconomic status. These data were collected using an adapted questionnaire from the second National Health and Family Survey, originally consisting of 29 questions to create a Standard of Living Index²⁶. Those questions relevant to the study population were used, with a total of 13 weighted questions for a maximum score of 26. Weights of items such as type of housing, community or household water and toilet facilities, household possession of televisions/radios, and ownership of livestock, were developed by the International Institute of Population Sciences in India and based on a priori knowledge of the significance of each indicator in determining household socioeconomic status²⁷.

Additional demographic information was collected from each participant, including religion, caste, and sources of income in the previous year. Time per day spent watching TV and information on health-seeking behaviour (type and frequency of healthcare sought) were collected. Religion was assessed as a binary variable (Hindu or 'other'), and caste was given the categories of low caste (scheduled tribes and caste), lower-middle caste (backwards castes, most backwards castes, other backwards castes), middle-upper caste (general category), and high caste (Brahmin caste). Data on occupation were collected by asking about sources of income within the last year, including local labour, agriculture, livestock, migrant work, merchant work, shop/small business, government schemes, or *Mahatma Gandhi National Rural Employment Guarantee Act* (MGNREGA). For ease of interpretation, livelihood information was given the categories of local labour, farming (agriculture and/or livestock), migrant work, business (merchant and/or small business owner), and government funds (MGNREGA and/or other government schemes) and assessed as binary variables ('yes' or 'no' for each individual source of income). Time per day spent watching television was collapsed into four categories: less than 0.5 hours per day, between 0.5 hours and 1 hour per day, more than 1 hour and up to 2 hours per day, or more than 2 hours per day.

Age was categorized into four groups – 20–34, 35–49, 50–64, and ≥ 65 years – based on similar methods used by Shrivastava et al¹⁶ and Murugesan et al¹³. Type of health care typically accessed included government, private, natural (ie ayurvedic and/or other alternative medicines), and none, and each was assessed as a binary variable. Frequency of healthcare visits was collapsed into three categories: once a month or more, less than once a month but more than once a year, and once a year or less. Information on school grade achieved was collected and had a range of 0–15, with anything greater than 12 indicating post-secondary education. For

ease of interpretation, this variable had the categories of no schooling, primary education (grades 1–8), secondary education (grades 9–11), and graduate or post-secondary education (grades ≥ 12), based on similar methods used previously among Indian populations^{13,24}.

Knowledge on diabetes was collected using a validated questionnaire developed by Mohan et al (2005)²⁴, and included questions about diabetes such as risk factors, complications, and prevention. The first question, 'Do you know what diabetes is?' acted as a screening question such that those who answered 'no' were automatically given a score of 0 and did not answer the remaining questions. For each response, weights were applied according to Mohan et al (2005) to calculate a composite knowledge score ranging from 0 to 8²⁴.

Statistical analysis

Data were cleaned using Microsoft Excel. All statistical analyses were conducted using Stata/IC v16.1 (StataCorp; <http://www.stata.com>). Due to zero inflation and heteroscedasticity of residuals, a linear regression model was not appropriate to model the composite knowledge score as a continuous variable. Therefore, diabetes knowledge had the categories of score of 0 (no knowledge), scores of 1–4 (moderate knowledge), and scores of 5–8 (good knowledge) for all statistical models.

Descriptive analyses were first conducted to establish the sociodemographic characteristics of the study population, overall diabetes knowledge scores, knowledge scores broken down by education level, and the proportion of the study population who answered questions correctly in the diabetes knowledge questionnaire. Following this, collinearity was tested by calculating the intraclass correlation coefficient. A coefficient higher than 0.8 was the cut-off point used to determine collinearity between variables. The linearity assumption was also tested against continuous predictor variables (such as age) and the log-likelihood of the outcome. Because the outcome variable contained three categories, a normal logistic regression could not be used to test linearity. Therefore, each predictor variable had to be tested against binary categories of the log-likelihood of the outcome (ie knowledge score in category 1 v 0, 2 v 0, and 1 v 2) using logistic regression. A LOWESS curve was first used to assess the linearity assumption, followed by the inclusion of a quadratic term. Variables that proved to be non-linear and could not be modelled with a quadratic term were then categorized.

To control for confounding bias, all individuals with self-reported diagnosed diabetes were omitted from the data analysis. All sociodemographic factors were then fit for univariable analysis with the outcome, using 'no knowledge' as the referent outcome category. These variables included age, sex, wealth index, education, religion, caste, methods of earning income, rural index, television exposure, and type and frequency of healthcare usually received. Only independent variables significant to a liberal p -value of 0.2 were short-listed for inclusion in the initial multivariable model.

Following this, a manual backward elimination method was used to fit an adjusted multinomial regression model to assess associations between sociodemographic factors and knowledge score categories, with a knowledge score of zero as the referent. Due to clustering of the data by village, village was added to the model as a random effect variable. To incorporate the use of a random effect in a multinomial model, the Stata program GLLAMM was used to fit a generalized linear latent and mixed model, using village as a discrete random effect, and knowledge score as a polytomous outcome with the *mlogit* link. Adaptive quadrature was also used to ensure the most precise estimates were given for the log-likelihood of the two-level model²⁸. Following the backward elimination process, each variable was removed independently from the full model and the partial model was then tested for significance ($p < 0.05$), using likelihood-ratio tests and Wald tests. As each variable was removed from the model, it was simultaneously tested for confounding by determining if any coefficients of interest changed by 20% or more following exclusion. All plausible interactions were then generated and assessed for significance using a p -value of < 0.05 . No confounding was found to be present, and no interaction variables were statistically significant in the adjusted multinomial model.

After the multinomial model was finalized, diagnostics were performed for GLLAMM using the *gllapred* command²⁸. Upper residuals (empirical Bayes predictions) were produced, along with Pearson and deviance residuals. Upper residuals were tested for homoscedasticity and normality, and Pearson and deviance residuals were used to assess outliers and their impact on the model. There were no outliers with a strong enough impact to justify their removal from the model, and no participants with a missing knowledge score. Therefore, only participants who did not fit the inclusion criteria for the model (ie were previously diagnosed with diabetes) were removed for analysis.

Ethics approval

This study received ethics clearance for research on human participants from the University of Guelph Research Ethics Board (certificate 12MY023). Permission for the study was granted by the High Commission of India in Ottawa, Canada. Upon arrival to the research site and prior to the recruitment process, researchers sought and received permission for the study from local authorities (panchayat councils, local police officials, and hospital medical staff). Informed verbal consent was obtained from all research participants prior to enrollment in the study.

Results

The response rate among the 812 recruited individuals was 92.7%, with 753 individuals completing the diabetes knowledge survey (341 males and 412 females). Descriptive characteristics of the study participants are presented in Table 1. The average age of participants was 47 years (standard deviation ± 14.7 years). Over half of the population (59.1%) had no formal schooling, with only 35.1% reporting full literacy (could read and write). Only about one-third of the population reported knowing about diabetes. More men reported having any knowledge of diabetes

when compared to women (36.9% v 30.3%); however, this difference was not significant. Those who reported being aware of diabetes were invited to participate in the full diabetes knowledge survey. Full results of the diabetes knowledge questionnaire are reported in Table 2.

Of those who reported awareness of diabetes, 62.6% answered 'yes' to the question, 'Do you know if diabetes is increasing?'. About half of the participants who were aware of diabetes believed that it is preventable. In total, only 16.4% and 17.0% achieved moderate and good knowledge scores, respectively. Even among those with the highest level of education, 42.2% did not know what diabetes was.

Literacy and education had high collinearity (intraclass correlation coefficient of 0.84), leading to the omission of literacy as a predictor variable. After individuals with self-reported diagnosed diabetes were removed, 705 individuals were included in statistical analyses. After assessing univariable analyses of each predictor variable, all variables were included for multivariable analysis except religion, television exposure, and migrant work as a source of income. The final multinomial regression model is presented in Table 3. Wealth, education, rurality, frequency of healthcare visits,

ownership of a business, and government funds as a source of income were all associated with moderate and/or good knowledge of diabetes in the final model. Notably, education was strongly associated with a good knowledge score in the final model. Specifically, those who were in the highest education category (having graduated high school or who had higher post-secondary education) were greater than 11 times more likely to have a good knowledge score over no knowledge, compared to those having no formal education as the referent (relative risk ratio (RRR) 11.1, 95% confidence interval (CI) 4.4, 27.6). A gradual increase in RRRs for having a good knowledge score was seen with each increasing category of education – primary (grades 1–8), secondary (grades 9–11), and high-school graduate or post-secondary (grades ≥ 12), respectively (RRR=3.2, 8.4, 11.1); however, this increasing trend was not seen for having a moderate knowledge score (RRRs= 1.8, 3.2, 1.3). Having a good knowledge score was also positively associated with frequency of healthcare usage, meaning those with less frequent healthcare visits had a lower relative risk of having a good knowledge score. Rurality was negatively associated with any knowledge of diabetes, indicating that those participants living in more rural locations had lower awareness and knowledge of diabetes.

Table 1: Descriptive characteristics of a sample of individuals in rural Tamil Nadu

Characteristic	Male (n=341, 45%) % or mean ± SD	Female (n=412, 55%) % or mean ± SD	Total (n=753) % or mean ± SD
Age (years)	48.1 ± 14.77	46.4 ± 14.61	47.14 ± 14.7
20–34	20.2	23.1	21.8
35–49	33.7	35.7	34.8
50–64	30.8	26.9	28.8
≥65	15.3	14.3	14.7
Wealth index (range 1–26)	11.7 ± 4.44	10.3 ± 4.76	10.9 ± 4.67
Low (1–9)	31.7	48.5	40.9
Middle (10–18)	60.1	44.7	51.7
High (19–26)	8.2	6.8	7.4
Grade standard achieved	4.1 ± 4.57	2.17 ± 3.76	3.04 ± 4.25
No schooling	45.6	71.6	59.1
Primary (grades 1–8)	33.4	17.1	24.5
Secondary (grades 9–11)	12.2	8.5	10.2
High-school graduate or post-secondary	8.9	2.8	5.5
Literate (can read and write)	49.3	23.3	35.1
Caste			
Lower caste (ST, SC)	11.3	16.7	14.1
Lower-middle caste (BC, MBC, OBC)	80.5	55.7	67.4
Middle-upper caste (GC)	5.0	10.2	7.7
Upper caste (Brahmin)	3.1	17.5	10.7
Self-reported methods of earning income in the last year			
Local day labour	49.4	58.5	54.4
Farming (agriculture and/or livestock)	69.5	54.1	61.1
Migrant work/labour	12.9	8.3	10.4
Local business and/or merchant work	16.7	12.9	14.6
Government funding (MGNREGA and/or government scheme)	19.1	35.7	28.2
Family history of diabetes	11.1	9.5	10.2
Diagnosed with diabetes	6.2	6.1	6.1

BC, Backward Class. GC, General Category. MBC, Middle Backward Class. MGNREGA, Mahatma Gandhi National Rural Employment Guarantee Act. OBC, Other Backward Class. SC, Scheduled Caste. ST, Scheduled Tribe.

Table 2: Questions correctly answered by a rural Tamil Nadu population from a diabetes knowledge survey

Question	n		% of total population
	n [†]	% [†]	
1. Know what diabetes is?	251		33.3
2. Know diabetes is increasing?	154	62.6	20.5
3. Know contributing factors of diabetes?			
Obesity	13	5.2	1.7
Decreased physical activity	27	10.8	3.4
Family history of diabetes	19	7.6	2.5
Mental stress	13	5.2	1.7
Consuming more sweets	124	50.4	16.5
4. Know diabetes can cause complications in organs?	51	20.6	6.8
5. Know diabetes can be prevented?	123	49.6	16.3

[†] Percentage of participants who knew of diabetes (ie responded 'yes' to first question).

Table 3: Associations of sociodemographic factors in the final multinomial logistic regression model on diabetes knowledge score in a rural Tamil Nadu population

Variable	Moderate knowledge (score 1–4)				Good knowledge (score 5–8)			
	RRR	95%CI		p-value	RRR	95%CI		p-value
Wealth index (1–26)	1.77	1.34	2.34	<0.0001	1.47	1.12	1.93	0.005
Education								
No education (ref)	–	–	–	–	–	–	–	–
Primary (grades 1–8)	1.77	1.02	3.08	0.043	3.21	1.72	5.99	<0.0001
Secondary (grades 9–11)	3.20	1.46	7.03	0.004	8.39	3.73	18.89	<0.0001
High-school graduate or post-secondary	1.32	0.40	4.34	0.649	11.07	4.44	27.61	<0.0001
Methods of earning income within the last year (yes/no)								
Business (merchant and/or shop owner)	2.18	1.14	4.14	0.018	2.12	1.10	4.06	0.024
Government funds (MGNREGA and/or government schemes)	3.27	1.93	5.54	<0.0001	2.39	1.31	4.36	0.005
Rural index [†]	0.34	0.19	0.59	<0.0001	0.43	0.24	0.74	0.003
Frequency of interactions with healthcare system								
Once a month or more (ref)	–	–	–	–	–	–	–	–
Less than once a month, but more than once a year	0.79	0.43	1.47	0.460	0.32	0.16	0.61	0.001
Once a year or less	0.73	0.37	1.44	0.364	0.30	0.15	0.61	0.001

[†] Positive scores reflect increasingly rural areas.

CI, confidence interval. MGNREGA, Mahatma Gandhi National Rural Employment Guarantee Act. RRR, relative risk ratio.

Discussion

This cross-sectional survey of adults residing in rural areas of Tamil Nadu showed poor knowledge of diabetes among the population. All participants – even subgroups with higher wealth and education – demonstrated a lack of knowledge of diabetes. Among those in the highest education category, over 40% had a knowledge score of zero, meaning they answered 'no' to the question, 'Do you know what diabetes is?'. Such results indicate that, even among the most educated individuals in the study population, general knowledge of diabetes was low. Although these findings are consistent with other studies^{12,14,17,24,29,30}, this rural region of Tamil Nadu demonstrated one of the lowest levels of knowledge in India reported to date, with 66% of participants being unaware of diabetes altogether. A study conducted by Shrivastava et al (2015) in another region of rural Tamil Nadu assessed knowledge and self-care practices among patients with diabetes and showed that, even among those with diabetes, knowledge of the disease was poor. Indeed, the study found that 49% of participants with diagnosed diabetes believed that diabetes was curable¹⁶. A similar qualitative study looking deeper into themes regarding diabetes knowledge among individuals with diabetes, showed that 96% of participants answered, 'I don't know' when asked, 'what happens inside your body when you get sick with diabetes?'³¹. The low awareness of diabetes shown in these studies corresponds with the present study's findings and should be considered a serious public health concern in rural regions of

South India.

Although levels of diabetes knowledge and awareness are consistently low across most studies in rural India, awareness varies across other regions in South Asia. A study conducted with adults from both urban and rural regions of Punjab, Pakistan, found that 86% of respondents had heard of diabetes³². The higher levels of education and socioeconomic status of this population could be an explanation for the heightened knowledge scores as compared to the present study. Another study conducted on a lower-middle class urban population in Chennai, the capital of the state of Tamil Nadu, showed that over 90% of the general population knew of diabetes¹³. Such findings suggest that diabetes awareness and knowledge may be better in urban regions, where information, messaging, and resources for diabetes may be more accessible. The association between higher wealth, education, and lower rurality and diabetes knowledge is consistent with other studies in India^{12,13,17}, and other low- and middle-income countries, including Jordan³³, Bangladesh³⁴, Southeast Ethiopia³⁵, Oman³⁶, and Pakistan³².

Doctor-patient interactions in healthcare settings are important opportunities for patient education, as shown by the strong association between frequency of healthcare visits and diabetes knowledge. This finding corresponds with previous research in the study region showing that patients with diabetes perceived doctors as the most important source of knowledge on diabetes³¹.

However, this study adds to the body of evidence suggesting that healthcare professionals in rural regions of South India do not adequately educate patients on diabetes; indeed, previous research in the study site³¹, and India in general^{37,38}, found that both public and private healthcare practitioners often failed to provide even basic health education and support to patients with non-communicable diseases.

An intriguing result was the association between source of income and knowledge score. Individuals working as a merchant or shop owner were more knowledgeable about diabetes, perhaps due to increased income and socioeconomic status, allowing for more exposure and access to accurate health information when compared to farmers or labourers. Individuals of higher socioeconomic status may also be at increased risk of cardiometabolic diseases such as diabetes, perhaps increasing the likelihood that a healthcare professional might educate them on such topics, or that they access information from peers of similar social status and diabetes risk³⁹. The nature of being a merchant and/or shop owner also creates opportunities for social interaction with community members, which may lead to discussions on health topics. Conversely, little exposure to people living with diabetes could also be a plausible explanation for the association between low diabetes knowledge and increased rurality in the present study, particularly given the lower prevalence of diabetes in rural versus urban and urbanizing settings⁹.

Accessing government schemes (eg MGNREGA) as a source of income was strongly associated with increased knowledge of diabetes. To the authors' knowledge, this association has not yet been established in any previous studies. The MGNREGA scheme provides employment security for adult (over age 18 years) applicants who reside in rural households throughout all districts of India⁴⁰. The main goal of the MGNREGA program is to provide employment opportunities within a radius of 5 km of a person's home for at least 100 days in a year. Most jobs involve manual unskilled labour and pay minimum wage; however, some higher skill jobs (eg project supervisor) are available with higher compensation⁴¹. MGNREGA creates work projects for members of rural communities involving different forms of labour, sometimes including the construction of permanent assets in participating communities, such as wells, roads, and bridges^{40,42}. While critiques of MGNREGA exist^{41,43-47}, involvement in this program throughout rural communities has the potential to foster local economic and employment activity, improve household income security, and improve quality of life^{41,42,48}.

While the association between diabetes knowledge and MGNREGA involvement has not been previously noted in the literature, this finding begets a number of plausible hypotheses that bear further exploration. Household income security through MGNREGA participation has been associated with increased household expenditure on education and health care^{41,42,49-51}, possibly increasing healthcare access, and thereby exposure to diabetes knowledge. However, MGNREGA wages are often delayed and unpredictable, and may be insufficient on their own to sustain households^{42,43,45,47,48,50}. Additionally, involvement in government

programs such as MGNREGA may improve participation in other government social welfare programs, although research in this area is lacking. One study in particular found that women involved in MGNREGA had a high awareness of other existing government schemes, with some even expressing concern for over-dependency on government benefits and programs⁴². Engagement in several government or social welfare programs may have the potential to foster trust and improve uptake in other government services and sources of information. In the present study, it is possible that those participants involved in MGNREGA were more likely to seek and trust health information from government sources. Lastly, MGNREGA can improve community cohesion and bonding among those involved in the program^{42,48}. Involvement in MGNREGA work may provide a platform to discuss common issues and interests, including health concerns⁴². Overall, the association between MGNREGA participation and diabetes knowledge is novel and possible mechanisms should be explored further.

Several studies highlight understandings and perceptions of diabetes among Indian populations that occasionally conflict with biomedical models. For example, a common perception in India is that consuming excess sugar is a primary cause of diabetes^{14,16,17,24,29}. Additionally, 'tension' or mental stress are often also cited as direct causes of diabetes^{14,17}, and herbal or religious remedies are often recognized as effective treatments for diabetes^{12,14}. Such patterns are consistent with the present study, as the most common perceived risk factor of diabetes was consuming sweets (16.5% of the study population). Correspondingly, as described in a previous study in the same region, the local colloquial term for diabetes was translated as 'sugar disease'³¹. Mental stress was also reported as a risk factor by 5% of those who knew of diabetes, the same proportion who reported obesity as a risk factor. This exemplifies how cultural and local understandings of health and disease (such as 'tension') may influence perceptions of diabetes causation^{14,31}. Evidence also indicates that, for many individuals in South Asia, family and friends are a main source of information on diabetes^{52,53}. Reliance on social networks for health information further perpetuates localized understandings of diabetes, grounded in experiences of individuals rather than information from health authorities.

The low number of individuals in this study who reported obesity as a risk factor to diabetes (1.7% of the total population) is particularly concerning, considering obesity is one of the strongest predictors of type 2 diabetes^{4,8-10}. Such findings correspond with a similar study investigating diabetes knowledge in a rural north-eastern Indian population, which found that only 40% of those who were overweight knew they had an increased risk of diabetes¹⁵. Sociocultural constructs regarding what is 'healthy' and 'unhealthy' may perpetuate this perception; indeed, some studies suggest that overweight and obesity are perceived as 'healthy' in subpopulations in rural India, especially among low socioeconomic status individuals, because overweight can be a sign of wealth and food security⁵².

The fragmented healthcare system in India, along with poor investments in public health initiatives and health education, limit

access to reputable and relevant information regarding health and disease, especially for rural populations. Many studies highlight the difficulty of receiving care for simple health issues, often citing the unavailability of doctors, long wait times, high costs, and lack of healthcare coordination^{54,55}. Despite the greater expense, private health care is often perceived as superior to public health care for major health problems such as diabetes⁵⁵. Regardless, both public and private healthcare facilities often face resource constraints that limit their ability to disseminate information regarding diabetes. Further, poverty and social inequities act as barriers to accessing health care in rural India. Healthcare centres are often located in urban cities (thus requiring transportation), are focused on tertiary care, and only affordable to the urban affluent, with rural poor individuals being faced with limited healthcare options⁵⁶. Many individuals in rural areas face financial hardships and use their income to sustain daily living, often avoiding healthcare services unless for life-threatening conditions⁵⁴.

India is currently grappling with an epidemiological transition that is driving an increasing burden of non-communicable diseases such as diabetes². So far, diabetes prevention efforts have been woefully unsatisfactory in India, especially in rural areas^{56,57}. It is therefore crucial and timely to improve efforts and allocate resources to alleviate the burden of diabetes. The associations of sociodemographic factors with diabetes knowledge in this study highlight priority areas for targeting initial public health efforts in Tamil Nadu. Specifically, efforts should emphasize the dissemination of accurate knowledge of diabetes signs, symptoms, prevention, and treatment to rural and isolated regions where populations lack access to formal education and seldomly interact with healthcare systems. Importantly, improved knowledge on diabetes has been associated with positive attitudes and better self-care practices towards diabetes treatment and prevention^{16,33,35,36}. Thus, investing in stronger public health efforts to improve healthcare access, quality, and focusing on non-communicable disease prevention and treatment, presents a crucial tool for lessening the severity and impacts of the diabetes epidemic in India. Additionally, structural factors grounded in economic and political realities – for example, food environments, access to sustainable livelihoods, and availability of recreational opportunities – are also crucial components to preventing and managing non-communicable diseases, and must be incorporated

into any regional or national strategy to prevent burdens of diabetes^{58,59}.

Strengths of this study include the use of systematic random sampling to reduce sampling bias, the use of validated tools to measure socioeconomic status and diabetes knowledge to reduce the risk of misclassification bias, and the use of a statistical modelling technique that accounted for confounding bias and incorporated a polytomous outcome. This study also examined rurality on a continuous scale instead of a binary urban–rural divide, allowing for increased granularity in examining the relationship between rurality and diabetes knowledge. This study also had some limitations. Importantly, cross-sectional surveys are unable to establish causation. Because most sociodemographic data and knowledge on diabetes was self-reported, some data may have been influenced by misreporting or social desirability bias. Finally, weaknesses of the diabetes knowledge questionnaire have been well documented²⁴, and include open-ended questions being inhibited by memory and recall bias, and closed-ended questions possibly encouraging respondents to provide guesses instead of informed answers.

Conclusion

The present study contributes to the large body of research regarding diabetes knowledge in India and South Asia. Specifically, this study sought to investigate the overall knowledge of diabetes in a rural population, as well as to examine the influence of different sociodemographic factors on knowledge of diabetes. Overall, this study highlighted the low levels of knowledge regarding diabetes in this rural population in Tamil Nadu. Positive associations were identified between knowledge score and wealth, education, MGNREGA and business ownership as sources of income, as well as frequency of health care received. Increasing rurality was negatively associated with knowledge score. Given the high prevalence of diabetes as well as prediabetes in this specific rural population⁹, as well as in the Tamil Nadu state in general⁷, this lack of general knowledge presents a major public health concern. Findings on the sociodemographic factors associated with knowledge of diabetes underscore the need for targeted education programming that aims to improve health literacy and diabetes awareness among those of lower socioeconomic status who lack formal education and reside in isolated and rural regions.

REFERENCES:

- 1 International Diabetes Foundation. *IDF diabetes atlas*. 9th edn. Brussels: International Diabetes Foundation, 2019.
- 2 Indian Council of Medical Research, Public Health Foundation of India, Institute for Health Metrics and Evaluation. *India: health of the nation's states. The India State-Level Disease Burden Initiative*. New Delhi: Indian Council of Medical Research, Public Health Foundation of India, Institute for Health Metrics and Evaluation, 2017.
- 3 Pradeepa R, Mohan V. Prevalence of type 2 diabetes and its complications in India and economic costs to the nation. *European Journal of Clinical Nutrition* 2017; **71**: 816-824. DOI link, PMID:28422124
- 4 Gupta SK, Singh Z, Purty AJ, Kar M, Vedapriya D, Mahajan P, et al. Diabetes prevalence and its risk factors in rural area of Tamil Nadu. *Indian Journal of Community Medicine* 2010; **35(3)**: 396-399. DOI link, PMID:21031104
- 5 Joshi SR. Diabetes care in India. *Annals of Global Health* 2015; **81(6)**: 830-838. DOI link, PMID:27108150
- 6 Nanditha A, Snehalatha C, Satheesh K, Susairaj P, Simon M, Vijaya L, et al. Secular TRends in DiabEtes in India (STRiDE-I):

- change in prevalence in 10 years among urban and rural populations in Tamil Nadu. *Diabetes Care* 2019; **42(3)**: 476-485. DOI link, PMID:30659076
- 7** India State-Level Disease Burden Initiative Diabetes Collaborators. 2018; **6(12)**: e1352-e1362. *The increasing burden of diabetes and variations among the states of India: The Global Burden of Disease Study*. Lancet Global Health , 1990-2016.
- 8** Anjana RM, Pradeepa R, Deepa M, Datta M, Sudha V, Unnikrishnan R, et al. Prevalence of diabetes and prediabetes (impaired fasting glucose and/or impaired glucose tolerance) in urban and rural India: phase I results of the Indian Council of Medical Research-India DIABetes (ICMR-INDIAB) study. *Diabetologia* 2011; **54(12)**: 3022-3027. DOI link, PMID:21959957
- 9** Little M, Humphries S, Patel K, Dodd W, Dewey C. Factors associated with glucose tolerance, pre-diabetes, and type 2 diabetes in a rural community of south India: a cross-sectional study. *Diabetology and Metabolic Syndrome* 2016; **8**: 21. DOI link, PMID:26958082
- 10** Oommen AM, Abraham VJ, George K, Jose VJ. Prevalence of risk factors for non-communicable diseases in rural and urban Tamil Nadu. *Indian Journal of Medical Research* 2016; **144(3)**: 460-471. DOI link, PMID:28139545
- 11** Muthunayanan L, Ramraj B, Russel JK. Prevalence of prediabetes and its associated risk factors among rural adults in Tamil Nadu. *Archives of Medicine and Health Sciences* 2015; **3**: 178-184. DOI link
- 12** Khan FN, Agne AA, Buttan S, Cherrington AL. Assessing diabetes knowledge and barriers to health services among patients with diabetes: a qualitative study in Delhi, India. *International Journal of Health Promotion and Education* 2013; **51(6)**: 290-299. DOI link
- 13** Murugesan N, Snehalatha C, Shobhana R, Roglic G, Ramachandran A. Awareness about diabetes and its complications in the general and diabetic population in a city in southern India. *Diabetes Research and Clinical Practice* 2007; **77(3)**: 433-437. DOI link, PMID:17291622
- 14** Mehta NV, Trivedi M, Maldonado LE, Saxena D, Humphries DL. Diabetes knowledge and self-efficacy among rural women in Gujarat, India. *Rural and Remote Health* 2016; **16(1)**: 3629. DOI link, PMID:26976745
- 15** Lau SL, Debarm R, Thomas N, Asha HS, Vasan KS, Alex RG, et al. Healthcare planning in north-east India: a survey on diabetes awareness, risk factors and health attitudes in a rural community. *Journal of the Association of Physicians of India* 2009; **57**: 305-309.
- 16** Shrivastava PS, Shrivastava SR, Ramasamy J. An epidemiological study to assess the knowledge and self care practices among type 2 diabetes mellitus patients residing in rural areas of Tamil Nadu. *Biology and Medicine* 2015; **53**: 002. DOI link
- 17** Deepa M, Bhansali A, Anjana RM, Pradeepa R, Joshi SR, Joshi PP, et al. Knowledge and awareness of diabetes in urban and rural India: The Indian Council of Medical Research India Diabetes Study (Phase I): Indian Council of Medical Research India Diabetes 4. *Indian Journal of Endocrinology and Metabolism* 2014; **18(3)**: 379-385. DOI link, PMID:24944935
- 18** Anjana RM, Deepa M, Pradeepa R, Mahanta J, Narain K, Das HK, et al. Prevalence of diabetes and prediabetes in 15 states of India: results from the ICMR-INDIAB population-based cross-sectional study. *Lancet Diabetes Endocrinology* 2017; **5(8)**: 585-596. DOI link
- 19** Taqi M, Bidhuri S, Sarkar S, Ahmad WS, Wangchok P. Rural healthcare infrastructural disparities in India: a critical analysis of availability and accessibility. *Journal of Multidisciplinary Research in Healthcare* 2017; **3(2)**: 125-149. DOI link
- 20** Banerjee S. Determinants of rural-urban differential in healthcare utilization among the elderly population in India. *BMC Public Health* 2021; **21**: 939. DOI link, PMID:34001026
- 21** Little M, Humphries S, Patel K, Dewey C. Factors associated with BMI, underweight, overweight, and obesity among adults in a population of rural south India: a cross-sectional study. *BMC Obesity* 2016; **3(1)**: 12. DOI link, PMID:26904203
- 22** Little M, Zivot C, Humphries S, Dodd W, Patel K, Dewey C. Burden and determinants of anemia in a rural population in south India: a cross-sectional study. *Anemia* 2018; **71(2)**: 3976. DOI link, PMID:30112198
- 23** Binson D, Catania JA. Random selection in a national telephone survey: a comparison of the Kish, next-birthday, and last-birthday methods. *Journal of Official Statistics* 2000; **16(1)**: 53.
- 24** Mohan D, Raj D, Shanthirani CS, Datta M, Unwin NC, Mohan V. Awareness and knowledge of diabetes in Chennai – The Chennai Urban Rural Epidemiology Study. *Journal of the Association of Physicians of India* 2005; **53**: 283-287.
- 25** Weinert C, Boik RJ. MSU Rurality Index: development and evaluation. *Research in Nursing and Health* 1995; **18(5)**: 453-464. DOI link, PMID:7676078
- 26** International Institute for Population Sciences. *National Family Health Survey (NFHS-2)*. 2007. Available: web link (Accessed 21 July 2020). Mumbai: International Institute for Population Sciences.
- 27** Ebrahim S, Kinra S, Bowen L, Andersen E, Ben-Shlomo Y, Lyngdoh T, et al. The effect of rural-to-urban migration on obesity and diabetes in India: a cross-sectional study. *PLoS Medicine* 2007; **7(4)**: e1000268. DOI link, PMID:20436961
- 28** Rabe-Hesketh S, Skrondal A, Pickles A. Reliable estimation of generalized linear mixed models using adaptive quadrature. *The Stata Journal* 2002; **2(1)**: 1-21. DOI link
- 29** Muninarayana C, Balachandra G, Hiremath SG, Iyengar K, Anil NS. Prevalence and awareness regarding diabetes mellitus in rural Tamaka, Kolar. *International Journal of Diabetes in Developing Countries* 2010; **30(1)**: 18-21. DOI link, PMID:20431801
- 30** Chavan GM, Waghachavare VB, Gore AD, Chavan VM, Dhobale RV, Dhumale GB. Knowledge about diabetes and relationship between compliance to the management among the diabetic patients from rural area of Sangli District, Maharashtra, India. *Journal of Family Medicine and Primary Care* 2015; **4(3)**: 439-443.
- 31** Little M, Humphries S, Patel K, Dewey C. Decoding the type 2

- diabetes epidemic in rural India. *Medical Anthropology* 2017; **26(2)**: 96-110. DOI link, PMID:27644458
- 32** Gillani A, Islam FA, Hayat K, Atif N, Yang C, Chang J, et al. Knowledge, attitudes and practices regarding diabetes in the general population: a cross-sectional study from Pakistan. *International Journal of Environmental Research and Public Health* 2018; **15(9)**: 1906. DOI link, PMID:30200534
- 33** Alsous M, Jalil MA, Odeh M, Al Kurdi R, Alnan M. Public knowledge, attitudes and practices toward diabetes mellitus: a cross sectional study from Jordan. *PLoS One* 2019; **14(3)**: e0214479. DOI link, PMID:30925187
- 34** Fottrell E, Ahmed N, Shaha SK, Jennings H, Kuddus A, Morrison J, et al. Diabetes knowledge and care practices among adults in rural Bangladesh: a cross-sectional survey. *British Medical Journal Global Health* 2018; **3(4)**. DOI link, PMID:30057800
- 35** Kassahun CW, Mekonen AG. Knowledge, attitude, practices and their associated factors towards diabetes mellitus among non-diabetes community members of Bale Zone administrative towns, South East Ethiopia. A cross-sectional study. *PLoS One* 2017; **12(2)**: e0170040. DOI link, PMID:28152066
- 36** Shafae MA, Al-Shukaili S, Rizvi SG, Farsi YA, Khan MA, Ganguly SS, et al. Knowledge and perceptions of diabetes in a semi-urban Omani population. *BioMed Central Public Health* 2008; **8(1)**. DOI link, PMID:18644163
- 37** Wangoo SK, Maji D, Das AK, Rao PV, Moses A, Sethi B, et al. Barriers and solutions to diabetes management: an Indian perspective. *Indian Journal of Endocrinology and Metabolism* 2013; **17(4)**: 594-601. DOI link, PMID:23961474
- 38** Hasan H, Zodpey S, Saraf A. Diabetologist's perspective on practice of evidence based diabetes management in India. *Diabetes Research and Clinical Practice* 2012; **95(2)**: 189-193. DOI link, PMID:22001282
- 39** Corsi DJ, Subramanian SV. Socioeconomic gradient and distribution of diabetes, hypertension, and obesity in India. *Cardiology: JAMA Network Open* 2019; **2(4)**: e190411. DOI link, PMID:30951154
- 40** Business Insider India Bureau. *What is Mgnrega and how can one enroll in the scheme?* 2019. Available: web link (Accessed 18 July 2020).
- 41** Sukhtankar S. *India's National Rural Employment Guarantee Scheme: what do we really know about the world's largest workfare program?* 2016. Available: web link (Accessed 9 August 2020).
- 42** Dandekar A, Bhandari V, Modi P. *An impact assessment study of the usefulness and sustainability of the assets created under Mahatma Gandhi National Rural Employment Guarantee Act (MGNREGA), in Sikkim.* 2010. Available: web link (Accessed 9 August 2020).
- 43** Dodd W, Wyngaarden S, Humphries S, Patel K, Majowicz S, Little M, et al. The relationship between MGNREGA and internal labour migration in Tamil Nadu, India. *European Journal of Development Research* 2018; **30(2)**: 178-194. DOI link
- 44** Dheeraj RP. A critical analysis of the Mahatma Gandhi National Rural Employment Guarantee Act (MGNREGA). *Global Journal of Interdisciplinary Social Sciences* 2017; **6(6)**: 8-12. DOI link
- 45** Nair M, Ariana P, Ohuma EO, Gray R, De Stavola B, Webster P. Effect of the Mahatma Gandhi National Rural Employment Guarantee Act (MGNREGA) on malnutrition of infants in Rajasthan, India: a mixed methods study. *PLoS One* 2013; **8(9)**: e75089. DOI link, PMID:24086447
- 46** Lakshminarayanan S. Role of government in public health: current scenario in India and future scope. *Journal of Family and Community Medicine* 2011; **18(1)**: 26-30. DOI link, PMID:21694957
- 47** Breitreuz R, Stanton C, Brady N, Pattison-Williams J, King ED, Mishra C, et al. The Mahatma Gandhi National Rural Employment Guarantee Scheme: a policy solution to rural poverty in India? *Development Policy Review* 2017; **35(3)**: 397-417. DOI link
- 48** Naikoo AA, Thakur SS. Impact of MGNREGA on people's health and quality of life among the people of Block Behibag of district Kulgam. *Asia Pacific Journal of Research* 2017; **1(4)**: 171-185.
- 49** Thomas M. *Guaranteeing children a better life? The impact of the Mahatma Gandhi National Rural Employment Guarantee Act on child health and learning in rural India.* Technical report, University of Texas at Austin. 2016. Available: web link (Accessed 10 September 2020).
- 50** Pandi J. Livelihood changes through MGNREGA: a study in Rural Tamil Nadu. *India Journal of Rural and Industrial Development* 2019; **7(1)**: 43-57.
- 51** Malangmeih L, Bhattacharyya K, Mitra A. Impact of MGNREGA on livelihood security of rural households: a case study in Bankura district of West Bengal State, India. *Economic Affairs* 2014; **59(2)**: 137-146. DOI link
- 52** Greenhalgh T, Helman C, Chowdhury AM. Health beliefs and folk models of diabetes in British Bangladeshis: a qualitative study. *British Medical Journal* 1998; **316**: 978-983. DOI link, PMID:9550958
- 53** Patel NR, Chew-Graham C, Bundy C, Kennedy A, Blickem C, Reeves D. Illness beliefs and the sociocultural context of diabetes self-management in British South Asians: a mixed methods study. *BioMed Central Family Practice* 2015; **16**: 58. DOI link, PMID:25958196
- 54** Bhojani U, Mishra A, Amruthavalli S, Devadasan N, Kolsteren P, De Henauw S, et al. Constraints faced by urban poor in managing diabetes care: patients' perspectives from South India. *Global Health Action* 2013; **6**: 22258. DOI link, PMID:24093885
- 55** Dodd W, King N, Humphries S, Little M, Dewey C. Self-reported morbidity and health service utilization in rural Tamil Nadu, India. *Social Science and Medicine* 2016; **161**: 118-125. DOI link, PMID:27285663
- 56** Siegel K, Narayan KM, Kinra S. Finding a policy solution to India's diabetes epidemic. *Health Affairs* 2008; **27(4)**: 1077-1090. DOI link, PMID:18607043
- 57** Pati MK, Swaroop N, Kar A, Aggarwal P, Jayanna K, Van Damme W. A narrative review of gaps in the provision of integrated care for noncommunicable diseases in India. *Public Health Reviews* 2020; **41(8)**. DOI link, PMID:32435518

58 Berkowitz SA, Karter AJ, Corbie-Smith G, Seligman HK, Ackroyd SA, Barnard LS, et al. Food insecurity, food "deserts," and glycemic control in patients with diabetes: a longitudinal analysis. *Diabetes Care* 2018; **41(6)**: 1188-1195. DOI link, PMID:29555650

59 Lumagbas LB, Coleman H, Bunders J, Pariente A, Belonje A, de Cock Buning T. Non-communicable diseases in Indian slums: re-framing the Social Determinants of Health. *Global Health Action* 2018; **11(1)**: 1438840. DOI link, PMID:29589512

This PDF has been produced for your convenience. Always refer to the live site <https://www.rrh.org.au/journal/article/6855> for the Version of Record.